

# Phosphorus Solubilizing Bacteria: Occurrence, Mechanisms and Their Role in Crop Production

Khosro Mohammadi

Department of Agronomy, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

**Abstract** Plants acquire phosphorus from soil solution as phosphate anion. It is the least mobile element in plant and soil contrary to other macronutrients. It precipitates in soil as orthophosphate or is absorbed by Fe and Al oxides through ligand exchange. Phosphorus solubilizing bacteria play role in phosphorus nutrition by enhancing its availability to plants through release from inorganic and organic soil P pools by solubilization and mineralization. Principal mechanism in soil for mineral phosphate solubilization is lowering of soil pH by microbial production of organic acids and mineralization of organic P by acid phosphatase. Use of phosphorus solubilizing bacteria as inoculants increases P uptake. These bacteria also increase prospects of using phosphatic rocks in crop production. Greater efficiency of P solubilizing bacteria has been shown through co-inoculation with other beneficial bacteria and mycorrhiza. This article incorporates the recent developments on microbial P solubilization into classical knowledge on the subject.

**Keywords** *Bacillus*, Mineralization, Organic Acids, Soil Phosphorus, Solubilization, *Pseudomonas*

## 1. Introduction

Phosphorus (P) is a major growth-limiting nutrient, and unlike the case for nitrogen, there is no large atmospheric source that can be made biologically available[1]. Root development, stalk and stem strength, flower and seed formation, crop maturity and production, N-fixation in legumes, crop quality, and resistance to plant diseases are the attributes associated with phosphorus nutrition. Although microbial inoculants are in use for improving soil fertility during the last century, however, a meager work has been reported on P solubilization compared to nitrogen fixation. Soil P dynamics is characterized by physicochemical (sorption - desorption) and biological (immobilization-mineralization) processes. Large amount of P applied as fertilizer enters in to the immobile pools through precipitation reaction with highly reactive  $Al^{3+}$  and  $Fe^{3+}$  in acidic, and  $Ca^{2+}$  in calcareous or normal soils[2].

Efficiency of P fertilizer throughout the world is around 10-25 % [3], and concentration of bio-available P in soil is very low reaching the level of  $1.0 \text{ mg kg}^{-1}$  soil [4]. Soil microorganisms play a key role in soil P dynamics and subsequent availability of phosphate to plants [5]. Inorganic forms of P are solubilized by a group of heterotrophic microorganisms excreting organic acids that dissolve phosphatic minerals and / or chelate cationic partners of the P ions directly,

releasing P into solution [6]. Phosphate solubilizing bacteria (PSB) are being used as biofertilizer since 1950s [7]. Release of P by PSB from insoluble and fixed / adsorbed forms is an important aspect regarding P availability in soils. There are strong evidences that soil bacteria are capable of transforming soil P to the forms available to plant. Microbial biomass assimilates soluble P, and prevents it from adsorption or fixation [8]. Microbial community influences soil fertility through soil microorganisms enhance the P availability to plants by mineralizing organic P in soil and by solubilizing precipitated phosphates [9, 10]. These bacteria in the presence of labile carbon serve as a sink for P by rapidly immobilizing it even in low P soils [11]. Subsequently, PSB become a source of P to plants upon its release from their cells. The PSB and plant growth promoting rhizobacteria (PGPR) together could reduce P fertilizer application by 50% without any significant reduction of crop yield [12, 13]. It infers that PSB inoculants / biofertilizers hold great prospects for sustaining crop production with optimized P fertilization.

## 2. Phosphorus Solubilizing Microorganisms

Evidence of naturally occurring rhizospheric phosphorus solubilizing microorganism (PSM) dates back to 1903 [14]. Bacteria are more effective in phosphorus solubilization than fungi [15]. Among the whole microbial population in soil, PSB constitute 1 to 50 %, while phosphorus solubilizing fungi (PSF) are only 0.1 to 0.5 % in P solubilization


\* Corresponding author:

khosromohammadi60@yahoo.com (Khosro Mohammadi)

Published online at <http://journal.sapub.org/re>

Copyright © 2012 Scientific & Academic Publishing. All Rights Reserved

potential[9]. Number of PSB among total PSM in north Iranian soil was around 88 %[16]. Microorganisms involved in phosphorus acquisition include mycorrhizal fungi and PSMs[17]. Among the soil bacterial communities, ectorhizospheric strains from *Pseudomonas* and *Bacilli*, and endosymbiotic rhizobia have been described as effective phosphate solubilizers. Strains from bacterial genera *Pseudomonas*, *Bacillus*, *Rhizobium* and *Enterobacter* along with *Penicillium* and *Aspergillus* fungi are the most powerful P solubilizers[18]. *Bacillus megaterium*, *B. circulans*, *B. subtilis*, *B. polymyxa*, *B. sircalmous*, *Pseudomonas striata*, and *Enterobacter* could be referred as the most important strains[19]. A nematofungus *Arthrobotrys oligospora* also has the ability to solubilize the phosphate rocks[20].



**Figure 1.** Ecological factors influencing the root exudation process and biofertilizers, adapted from Nihorimbere et al. (2011)

### 3. Occurrence of PSB

High proportion of PSM is concentrated in the rhizosphere, and they are metabolically more active than from other sources. Usually, one gram of fertile soil contains 10<sup>1</sup> to 10<sup>10</sup> bacteria, and their live weight may exceed 2,000 kg ha<sup>-1</sup>. Soil bacteria are in cocci (sphere, 0.5 μm), bacilli (rod, 0.5-0.3 μm) or spiral (1-100 μm) shapes. Bacilli are common in soil, whereas spirilli are very rare in natural environments[21]. The PSB are ubiquitous with variation in forms and population in different soils. Population of PSB depends on different soil properties (physical and chemical properties, organic matter, and P content) and cultural activities. Larger populations of PSB are found in agricultural and rangeland soils[22]. In north of Iran, the PSB count ranged from 0 to

10<sup>7</sup> cells g<sup>-1</sup> soil, with 3.98 % population of PSB among total bacteria[16]. Plant species, plant developmental stage and soil type have thus been indicated as major factors determining the composition of biofertilizers. That said, the extent to which the above-cited factors contribute to microbial communities is not fully understood and there are several contrasting reports in the literature indicating either plant or soil type as dominant factor. Owing to the above statement, it can be generalized that the diversity and predominance of rhizosphere microbial population depend on a number of abiotic and biotic factors prevailing in that particular ecological niche (Fig. 2).

### 4. Mechanisms of PSB

Some bacterial species have mineralization and solubilization potential for organic and inorganic phosphorus, respectively[23]. Phosphorus solubilizing activity is determined by the ability of microbes to release metabolites such as organic acids, which through their hydroxyl and carboxyl groups chelate the cation bound to phosphate, the latter being converted to soluble forms[24]. Phosphate solubilization takes place through various microbial processes / mechanisms including organic acid production and proton extrusion[25]. A wide range of microbial P solubilization mechanisms exist in nature and much of the global cycling of insoluble organic and inorganic soil phosphates is attributed to bacteria and fungi. Phosphorus solubilization is carried out by a large number of saprophytic bacteria and fungi acting on sparingly soluble soil phosphates, mainly by chelation-mediated mechanisms[18]. Inorganic P is solubilized by the action of organic and inorganic acids secreted by PSB in which hydroxyl and carboxyl groups of acids chelate cations (Al, Fe, Ca) and decrease the pH in basic soils[26]. The PSB dissolve the soil P through production of low molecular weight organic acids mainly gluconic and keto gluconic acids[27], in addition to lowering the pH of rhizosphere. The pH of rhizosphere is lowered through biotical production of proton/bicarbonate release (anion/cation balance) and gaseous (O<sub>2</sub>/CO<sub>2</sub>) exchanges. Phosphorus solubilization ability of PSB has direct correlation with pH of the medium.

Release of root exudates such as organic ligands can also alter the concentration of P in the soil solution. Organic acids produced by PSB solubilize insoluble phosphates by lowering the pH, chelation of cations and competing with phosphate for adsorption sites in the soil[25]. Inorganic acids e.g. hydrochloric acid can also solubilize phosphate but they are less effective compared to organic acids at the same pH. In certain cases phosphate solubilization is induced by phosphate starvation[28].

#### 4.1. Solubilization of Ca-bound P

Soil phosphates mainly the apatites and metabolites of phosphatic fertilizers are fixed in the form of calcium phosphates under alkaline conditions. Many of the calcium

phosphates, including rock phosphate ores (fluoroapatite, francolite), are insoluble in soil with respect to the release of inorganic P (Pi) at rates necessary to support agronomic levels of plant growth. Gerretsen[29] first showed that pure cultures of soil bacteria could increase the P nutrition of plants through increased solubility of Ca-phosphates. Their solubility increases with a decrease of soil pH. Phosphate solubilization is the result of combined effect of pH decrease and organic acids production[17]. Microorganisms through secretion of different types of organic acids e.g. carboxylic acid[27] and rhizospheric pH lowering mechanisms dissociate the bound forms of phosphate like  $\text{Ca}_3(\text{PO}_4)_2$ . Nevertheless, buffering capacity of the medium reduce the effectiveness of PSB in releasing P from tricalcium phosphates[30]. Acidification of the microbial cell surroundings releases P from apatite by proton substitution / excretion of  $\text{H}^+$  (accompanying greater absorption of cation than anions) or release of  $\text{Ca}^{2+}$ [31]. While, the reverse occurs when uptake of anions exceeds that of cation, with excretion of  $\text{OH}^-$  /  $\text{HCO}_3^-$  exceeding that of  $\text{H}^+$ . Carboxylic anions produced by PSB, have high affinity to calcium, solubilize more phosphorus than acidification alone[32].

Complexing of cations is an important mechanism in P solubilization if the organic acid structure favors complexation. It is controlled by nutritional, physiological and growth conditions of the microbial culture, but it is mostly due to the lowering of pH alone by organic acids or production of microbial metabolites[33]. Organic anions and associated protons are effective in solubilizing precipitated forms of soil P (e.g. Fe and Al - P in acid soils, Ca - P in alkaline soils), chelating metal ions that may be associated with complexed forms of P or may facilitate the release of adsorbed P through ligand exchange reactions. Calcium phosphate (Ca-P) release results from the combined effects of pH decrease and carboxylic acids synthesis, but proton release cannot be the single mechanism[27].

#### 4.2. Solubilization of Al/Fe bound P

Solubilization of Fe and Al occurs via proton release by PSB by decreasing the negative charge of adsorbing surfaces to facilitate the sorption of negatively charged P ions. Proton release can also decrease P sorption upon acidification which increases  $\text{H}_2\text{PO}_4^-$  in relation to  $\text{HPO}_4^{2-}$  having higher affinity to reactive soil surfaces. Carboxylic acids mainly solubilize Al-P and Fe-P[14] through direct dissolution of mineral phosphate as a result of anion exchange of  $\text{PO}_4^{3-}$  by acid anion, or by chelation of both Fe and Al ions associated with phosphate[34]. It is through root colonizing *pseudomonads* with high-affinity iron uptake system based on the release of  $\text{Fe}^{3+}$  chelating molecules i.e. siderophores. Moreover, carboxylic anions replace phosphate from sorption complexes by ligand exchange and chelate both Fe and Al ions associated with phosphate, releasing phosphate available for plant uptake after transformation. Ability of organic acids to chelate metal cations is greatly influenced by its molecular structure, particularly by the number of carboxyl and hy-

droxyl groups. Type and position of the ligand in addition to acid strength determine its effectiveness in the solubilization process. Phosphorus desorption potential of different carboxylic anions lowers with decrease in stability constants of Fe or Al organic acid complexes ( $\log K_{\text{Al}}$  or  $\log K_{\text{Fe}}$ ) in the order: citrate > oxalate > malonate / malate > tartrate > lactate > gluconate > acetate > formiate[35].

#### 4.3. Mineralization of organic P

Mineralization of soil organic P (Po) plays an imperative role in phosphorus cycling of a farming system. Organic P may constitute 4-90 % of the total soil P. Almost half of the microorganisms in soil and plant roots possess P mineralization potential under the action of phosphatases[36]. Alkaline and acid phosphatases use organic phosphate as a substrate to convert it into inorganic form[37]. Principal mechanism for mineralization of soil organic P is the production of acid phosphatases. Release of organic anions, and production of siderophores and acid phosphatase by plant roots / microbes or alkaline phosphatase enzymes hydrolyze the soil organic P or split P from organic residues. The largest portion of extracellular soil phosphatases is derived from the microbial population[38]. *Enterobacter agglomerans* solubilizes hydroxyapatite and hydrolyze the organic P[39]. Mixed cultures of PSMs (*Bacillus*, *Streptomyces*, *Pseudomonas* etc.) are most effective in mineralizing organic phosphate.

### 5. Interaction of PSB with other Microorganism

Symbiotic relationship between PSB and plants is synergistic in nature as bacteriaprovide soluble phosphate and plants supply root borne carbon compounds (mainly sugars), that can be metabolized for bacterial growth. The PSM along with other beneficial rhizospheric microflora enhance crop production. Simultaneous application of *Rhizobium* with PSM or arbuscular mycorrhizae (AM) fungi has been shown to stimulate plant growth more than with their sole inoculation in certain situations when the soil is P deficient. Synergistic interactions on plant growth have been observed by coinoculation of PSB with  $\text{N}_2$  fixers such as *Azospirillum* and *Azotobacter*, or with vesicular arbuscular mycorrhizae [39, 40].

### 6. Amount of Solubilized P

The PSB solubilize the fixed soil P and applied phosphates resulting in higher crop yields[41]. Direct application of phosphate rock is often ineffective in the short time period of most annual crops. Acid producing microorganisms are able to enhance the solubilization of phosphatic rock. The PSB strains exhibit inorganic P-solubilizing abilities ranging between 25–42  $\mu\text{g P mL}^{-1}$  and organic P mineralizing abilities between 8–18  $\mu\text{g P mL}^{-1}$ [42]. The PSB in conjunction

with single super phosphate and rock phosphate reduce the P dose by 25 and 50 %, respectively[43]. *Pseudomonas putida*, *P. fluorescens* Chao and *P. fluorescens* Tabriz released 51, 29 and 62 % P, respectively; with highest value of 0.74 mg P / 50 mL from Fe<sub>2</sub>O<sub>3</sub>[44]. *Pseudomonas striata* and *Bacillus polymyxa* solubilized 156 and 116 mg P L<sup>-1</sup>, respectively (Rodríguez and Fraga, 1999). *Pseudomonas fluorescens* solubilized 100 mg P L<sup>-1</sup> containing Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub> or 92 and 51 mg P L<sup>-1</sup> containing AlPO<sub>4</sub> and FePO<sub>4</sub>, respectively[45].

## 7. Effect of PSB on Crop Production

Phosphate rock minerals are often too insoluble to provide sufficient P for crop uptake. Use of PSMs can increase crop yields up to 70 percent. Combined inoculation of arbuscular mycorrhiza and PSB give better uptake of both native P from the soil and P coming from the phosphatic rock[46]. Higher crop yields result from solubilization of fixed soil P and applied phosphates by PSB[40, 49]. Microorganisms with phosphate solubilizing potential increase the availability of soluble phosphate and enhance the plant growth by improving biological nitrogen fixation. *Pseudomonas* spp. enhanced the number of nodules, dry weight of nodules, yield components, grain yield, nutrient availability and uptake in soybean crop[47, 49]. Phosphate solubilizing bacteria

enhanced the seedling length of *Cicer arietinum*, while co-inoculation of PSM and PGPR reduced P application by 50 % without affecting corn yield[13]. Inoculation with PSB increased sugarcane yield by 12.6 percent[43]. Sole application of bacteria increased the biological yield, while the application of the same bacteria along with mycorrhizae achieved the maximum grain weight[48]. Single and dual inoculation along with P fertilizer was 30-40 % better than P fertilizer alone for improving grain yield of wheat, and dual inoculation without P fertilizer improved grain yield up to 20 % against sole P fertilization. Mycorrhiza along with *Pseudomonas putida* increased leaf chlorophyll content in barley[48]. Findings of Mohammadi[41] showed that inoculation of biofertilizers (PSB+ *Trichoderma* fungi) + application of FYM had a great influence on canola growth, height and grain yield in compared to control treatment (Fig. 2). Rhizospheric microorganisms can interact positively in promoting plant growth, as well as N and P uptake. Seed yield of green gram was enhanced by 24 % following triple inoculation of *Bradyrhizobium* + *Glomus fasciculatum* + *Bacillus subtilis*[50]. Growth and phosphorus content in two alpine *Carex* species increased by inoculation with *Pseudomonas fortinii*[51]. Integration of half dose of NP fertilizer with biofertilizer gives crop yield as with full rate of fertilizer; and through reduced use of fertilizers the production cost is minimized and the net return maximized.



Figure 2. Response of canola growth to biofertilizers (Mohammadi, 2011)

## 8. Conclusions

Soil P precipitated as orthophosphate and adsorbed by Fe and Al oxides is likely to become bio-available by bacteria through their organic acid production and acid phosphatase secretion. Although, high buffering capacity of soil reduces the effectiveness of PSB in releasing P from bound phosphates; however, enhancing microbial activity through P solubilizing inoculants may contribute considerably in plant P uptake. Phosphorus solubilizing bacteria mainly *Bacillus*, *Pseudomonas* and *Enterobacter* are very effective for increasing the plant available P in soil as well as the growth and yield of crops. So, exploitation of phosphate solubilizing bacteria through biofertilization has enormous potential for making use of ever increasing fixed P in the soil, and natural reserves of phosphate rocks.

## REFERENCES

- [1] Ezawa, T., S. E. Smith and F. A. Smith. 2002. P metabolism and transport in AM fungi. *Plant Soil* 244:221-230.
- [2] Hao, X., C. M. Cho, G. J. Racz and C. Chang. 2002. Chemical retardation of phosphate diffusion in an acid soil as affected by liming. *Nutr. Cycl. Agroecosys.* 64:213-224.
- [3] Isherword, K. F. 1998. Fertilizer use and environment. In: N. Ahmed and A. Hamid (eds.), *Proc. Symp. Plant Nutrition Management for Sustainable Agricultural Growth*. NFDC, Islamabad. pp. 57-76.
- [4] Goldstein, A. H. 2000. Bioprocessing of rock phosphate ore: essential technical considerations for the development of a successful commercial technology. *Proc. 4th Int. Fert. Assoc. Tech. Conf.* IFA, Paris. p. 220.
- [5] Richardson, A. E. 2001. Prospects for using soil microorganisms to improve the acquisition of phosphorus by plants. *Aust. J. Plant Physiol.* 28:897-906.
- [6] He, Z. L., W. Bian and J. Zhu. 2002. Screening and identification of microorganisms capable of utilizing phosphate adsorbed by goethite. *Comm. Soil Sci. Plant Anal.* 33:647-663.
- [7] Krasilnikov, N. A. 1957. On the role of soil micro-organism in plant nutrition. *Microbiologiya* 26:659-72.
- [8] Khan, K. S. and R. G. Joergensen. 2009. Changes in microbial biomass and P fractions in biogenic household waste compost amended with inorganic P fertilizers. *Bioresour. Technol.* 100:303-309.
- [9] Chen, Y. P., P. D. Rekha, A. B. Arunshen, W. A. Lai and C. C. Young. 2006. Phosphate solubilizing bacteria from subtropical soil and their tricalcium phosphate solubilizing abilities. *Appl. Soil Ecol.* 34:33-41.
- [10] Pradhan, N. and L. B. Sukla. 2005. Solubilization of inorganic phosphate by fungi isolated from agriculture soil. *African J. Biotechnol.* 5:850-854.
- [11] Bünemann, E. K., D. A. Bossio, P. C. Smithson, E. Frossard and A. Oberson. 2004. Microbial community composition and substrate use in a highly weathered soil as affected by crop rotation and P fertilization. *Soil Biol. Biochem.* 36:889-901.
- [12] Jilani, G., A. Akram, R. M. Ali, F. Y. Hafeez, I. H. Shamsi, A. N. Chaudhry and A. G. Chaudhry. 2007. Enhancing crop growth, nutrients availability, economics and beneficial rhizosphere microflora through organic and biofertilizers. *Ann. Microbiol.* 57:177-183.
- [13] Yazdani M., M. A. Bahmanyar, H. Pirdashti and M. A. Esmaili. 2009. Effect of Phosphate solubilization microorganisms (PSM) and plant growth promoting rhizobacteria (PGPR) on yield and yield components of Corn (*Zea mays* L.). *Proc. World Acad. Science, Eng. Technol.* 37:90-92.
- [14] Khan, M. S., A. Zaidi and P. A. Wani. 2007. Role of phosphate-solubilizing microorganisms in sustainable agriculture - A review. *Agron. Sustain. Dev.* 27:29-43.
- [15] Afzal, A. and A. Bano. 2008. Rhizobium and phosphate solubilizing bacteria improve the yield and phosphorus uptake in wheat (*Triticum aestivum* L.). *Int. J. Agri. Biol.* 10:85-88.
- [16] Fallah, A. 2006. Abundance and distribution of phosphate solubilizing bacteria and fungi in some soil samples from north of Iran. 18th World Congress of Soil Science, July 9-15, 2006, Philadelphia, Pennsylvania, USA.
- [17] Fankem, H., D. Nwaga, A. Deubel, L. Dieng, W. Merbach and F. X. Etoa. 2006. Occurrence and functioning of phosphate solubilizing microorganisms from oil palm tree (*Elaeis guineensis*) rhizosphere in Cameroon. *African J. Biotech.* 5:2450-2460.
- [18] Whitelaw, M. A. 2000. Growth promotion of plants inoculated with phosphate solubilizing fungi. *Adv. Agron.* 69:99-151.
- [19] Kucey, R. M. N., H. H. Janzen and M. E. Legget. 1989. Microbial mediated increases in plant available phosphorus. *Adv. Agron.* 42:199-228.
- [20] Duponnois, R., M. Kisa and C. Plenchette. 2006. Phosphate solubilizing potential of the nematofungus *Arthrobotrys oligospora*. *J. Plant Nutr. Soil Sci.* 169:280-282.
- [21] Baudoin E., E. Benizri and A. Guckert. 2002. Impact of growth stages on bacterial community structure along maize roots by metabolic and genetic fingerprinting. *Appl. Soil Ecol.* 19:135-145.
- [22] Yahya, A. and S. K. A. Azawi. 1998. Occurrence of phosphate solubilizing bacteria in some Iranian soils. *Plant Soil* 117:135-141.
- [23] Khiari, L. and L. E. Parent. 2005. Phosphorus transformations in acid light-textured soils treated with dry swine manure. *Can. J. Soil Sci.* 85:75-87.
- [24] Sagoe, C. I., T. Ando, K. Kouno and T. Nagaoka. 1998. Relative importance of protons and solution calcium concentration in phosphate rock dissolution by organic acids. *Soil Sci. Plant Nutr.* 44:617-625.
- [25] Nahas, E. 1996. Factors determining rock phosphate solubilization by microorganism isolated from soil. *World J. Microb. Biotechnol.* 12:18-23.
- [26] Stevenson, F. J. 2005. *Cycles of Soil: Carbon, Nitrogen, Phosphorus, Sulfur, Micronutrients*. John Wiley and Sons,

New York.

- [27] Deubel, A., Gransee and W. Merbach. 2000. Transformation of organic rhizodeposits by rhizoplane bacteria and its influence on the availability of tertiary calcium phosphate. *J. Plant Nutr. Soil Sci.* 163:387-392.
- [28] Gyaneshwar, P., G. N. Kumar, L. J. Parekh and P. S. Poole. 2002. Role of soil microorganisms in improving P nutrition of plants. *Plant Soil* 245:83-93.
- [29] Gerretsen, F. C. 1948. The influence of microorganisms on the phosphate intake by the plant. *Plant Soil* 1:51-81.
- [30] Stephen, J. and M. S. Jisha. 2009. Buffering reduces phosphate solubilizing ability of selected strains of bacteria. *World J. Agric. Sci.* 5:135-137.
- [31] Villegas, J. and J. A. Fortin. 2002. Phosphorus solubilization and pH changes as a result of the interactions between soil bacteria and arbuscular mycorrhizal fungi on a medium containing NO<sub>3</sub> as nitrogen source. *Can. J. Bot.* 80:571-576.
- [32] Staunton, S. and F. Leprince. 1996. Effect of pH and some organic anions on the solubility of soil phosphate: implications for P bioavailability. *Eur. J. Soil Sci.* 47:231-239.
- [33] Abd-alla, M. H. 1994. Phosphatases and the utilization of organic P by *Rhizobium leguminosarum* biovar viceae. *Lett. Appl. Microbiol.* 18:294-296.
- [34] Omar, S. A. 1998. The role of rock-phosphate-solubilizing fungi and vesicular-arbuscular mycorrhiza (VAM) in growth of wheat plants fertilized with rock phosphate. *World J. Microbiol. Biotechnol.* 14:211-218.
- [35] Ryan, P. R., E. Delhaize and D. L. Jones. 2001. Function and mechanism of organic anion exudation from plant roots. *Annl. Rev. Plant Physiol. Plant Mol. Biol.* 52:527-560.
- [36] Tarafdar, J. C. and N. Claasen. 1988. Organic phosphorus compounds as a phosphorus source for higher plants through the activity of phosphatases produced by plant roots and microorganisms. *Biol. Fert. Soils* 5:308-312.
- [37] Beech, I. B., M. Paiva, M. Caus and C. Coutinho. 2001. Enzymatic activity and within biofilms of sulphate-reducing bacteria. In: P. G. Gilbert, D. Allison, M. Brading, J. Verran and J. Walker (eds.), *Biofilm Community Interactions: chance or necessity?* BioLine, Cardiff, UK. pp. 231-239.
- [38] Dodor, D. E. and A. M. Tabatabai. 2003. Effect of cropping systems on phosphatases in soils. *J. Plant Nutr. Soil Sci.* 166:7-13.
- [39] Kim, K. Y., D. Jordan and G. A. McDonald. 1998. Effect of phosphate-solubilizing bacteria and vesicular-arbuscular mycorrhizae on tomato growth and soil microbial activity. *Biol. Fert. Soils* 26:79-87.
- [40] Zaidi, A., M. S., Khan and M. Amil. 2003. Interactive effect of rhizotrophic microorganisms on yield and nutrient uptake of chickpea (*Cicer arietinum* L.). *Eur. J. Agron.* 19:15-21.
- [41] Mohammadi, K., Ghalavand, A., Aghaalikhani, M., Heidari, G.R., Sohrabi, Y. 2011. Introducing the sustainable soil fertility system for chickpea (*Cicer arietinum* L.). *Afr J. Biotech.* 10(32): 6011-6020.
- [42] Tao, G., S. Tian, M. Cai and G. Xie 2008. Phosphate solubilizing and mineralizing abilities of bacteria isolated from soils. *Pedosphere.* 18:515-523.
- [43] Sundara, B., V. Natarajan and K. Hari. 2002. Influence of phosphorus solubilizing bacteria on the changes in soil available phosphorus and sugarcane yields. *Field Crops Res.* 77:43-49.
- [44] Goenadi, D. H., Siswanto and Y. Sugiarto. 2000. Bioactivation of poorly soluble phosphate rocks with a phosphorus-solubilizing fungus. *Soil Sci. Soc. Am. J.* 64:927-932.
- [45] Henri, F., N. N. Laurette, D. Annette, Q. John, M. Wolfgang, E. François-Xavier and N. Dieudonné. 2008. Solubilization of inorganic phosphates and plant growth promotion by strains of *Pseudomonas fluorescens* isolated from acidic soils of Cameroon. *African J. Microbiol. Res.* 2:171-178.
- [46] Cabello, M., G. Irrazabal, A. M. Bucsinszky, M. Saparrat and S. Schalamuck. 2005. Effect of an arbuscular mycorrhizal fungus, *G. mosseae* and a rock-phosphate-solubilizing fungus, *P. thomii* in *Mentha piperita* growth in a soilless medium. *J. Basic Microbiol.* 45:182-189.
- [47] Son, T. T. N., C. N. Diep and T. T. M. Giang. 2006. Effect of bradyrhizobia and phosphate solubilizing bacteria application on Soybean in rotational system in the Mekong delta. *Omo-nrice.* 14:48-57.
- [48] Mehrvarz, S., M. R. Chaichi and H. A. Alikhani. 2008. Effects of phosphate solubilizing microorganisms and phosphorus chemical fertilizer on yield and yield components of Barely (*Hordeum vulgare* L.). *Am-Euras. J. Agric. & Environ. Sci.* 3:822-828.
- [49] Mohammadi, K. 2011. Soil, plant and microbe interaction. Lambert Academic Publication. 120 pp.
- [50] Zaidi, A. and M. S. Khan. 2006. Co-inoculation effects of phosphate solubilizing microorganisms and *Glomus fasciculatum* on green gram - *Bradyrhizobium* symbiosis. *Turk. J. Agric.* 30:223-230.